FEP

Reference Scenarios Development based on Interaction Matrix and FEP for the Near-surface LILW Repository

, ,

150

FEP

6 가 .

Abstract

Systematic development procedure of radionuclide release scenarios was established based on FEP list and Interaction Matrix for near-surface LILW repository. FEPs were screened by experts' review in terms of domestic situation and combined into scenarios on the basis of Interaction Matrix analysis. Under the assumption of design scenario, system domain was divided into three sections: Near-field, Far-field and Biosphere. Subscenarios for each section were developed, and then scenarios for entire system were composed of sub-scenarios from each section. Finally, 6 reference scenarios for near-surface repository were selected through scenario screening.

1.

FEP(Features, Events, and Process) (Interaction Matrix) FEP 가 가 FEP 가 가 FEP 가 가 가 2. 가 가 가 , 2) 1) , 3) 가 4) , 5) 300 (100 200), 6) 가 [1]. ISAM FEP [1] 가 FEP . FEP FEP 2^n 가 가 n 가 가 FEP 1 FEP 2 Near-field, Far-field, Biosphere

FEP

FEP

2.1 Near-field

				, backfill, vault		Near-
field					가	
11010	N C . 1.1			•		
	Near-field					
•						
(1)	$(Normal\ Evolution) - NSS1$					
가	,	가				
						far-
C .14						
field	•					
(2)	(Colloid Transport) – NSS2					
backfill				(pseudo-colloid)		
	[2].					
(3) 가	(Gas Generation) – NSS3					
(3) 7 1	(Sub Generation) 1499					
				가		
	가					[2].
2.2 Far-field	d					
				Far-field		
		,	٦L	Tur field		
	•		가			
Far-field	가					
	Far-field					
(1) Fracture F	Flow – FSS1					
(2) Porous Flo	ow – FSS2					

フ	가
	가 .
(2)	(C. H.: LT
(3)	(Colloid Transport) – FSS3 フト
Near-field	71
(4)	(Undetected Features) – FSS4
	,
	•
2.3 Bioshpe	
2.5 Diosupe	
	, , , ,
	. 가
	가 .
·	
(1)	Water Resource)– BSS1
	•
(2)	(Discharge to Surface Water Body)– BSS2
` '	가 (, , ,
)	
(2)	(G. 'I I.G. I'). DCC2
(3)	(Soil and Sediment)- BSS3

.

3. 가 가 BSS1, BSS2, BSS3 Near-field Far-가 field 가 , 가 FEP 3 3.1 1: (1) **DS1-1** 가 가 porous flow (2) **DS1-2**

3.2 2:

(1) **DS2-1**

			가			
				•		
(2) DS2-2						
backfill				(psei	udo-colloid)	
			가			
()					
3.3	3 : 가					
(1) DS3-1						
					가	
	. Far-field			porous	flow	
·						
(2) DS3-2						
					가	
		()		
4.						
			EED			
			FEP	6	FEP	•
						Near-field, Far-field,
Biosphere						near-neid, Far-neid,

backfill

(pseudo-colloid)


가

	ISAM, the International Programme for Improving Long Term Safety Surface Radioactive Waste Disposal Facilities: Vault Safety Case Repo	_
[2] KINS, 1997		, KINS/HR-174, ,
[3] ,	, , 1994	,
[4] ,	, , 1995	가,


[5] SKB, Scenario Development Methodologies, SKB Technical Report 94-28, 1994


[6] SKB, The joint SKI/SKB scenario development project, SKB Technical Report 89-35, 1989

		1.	
	DS1-1 (NSS1+FSS2)	가	가 porous flow
	DS1-2 (NSS1+FSS4)	가 ,	가 ()
	DS2-1 (NSS2+FSS3)	backfill (pseudo-colloid) 가	
	DS2-2 (NSS2+FSS4)	backfill (pseudo-colloid) フト ()	
가	DS3-1 (NSS3+FSS2)	フト . Far-field	porous flow
	DS3-2 (NSS3+FSS4)	가	



1. 12×12





3. DS1-1